















- [14] N. Khuriyati, A. C. Sukartiko, and R. N. Alfiani, "Non-destructive measurement of antioxidant activity and water content in chili powder (*Capsicum annum* L.) using near-infrared spectroscopy," *Int. Food Res. J.*, vol. 29, no. 2, pp. 320–327, 2022.
- [15] F. W. Hong and K. S. Chia, "A review on recent near infrared spectroscopic measurement setups and their challenges," *Measurement*, vol. 171, p. 108732, 2021, doi: 10.1016/j.measurement.2020.108732.
- [16] K. D. C. Perera, G. K. Weragoda, R. Haputhanthri, and S. K. Rodrigo, "Study of concentration dependent curcumin interaction with serum biomolecules using ATR-FTIR spectroscopy combined with Principal Component Analysis (PCA) and Partial Least Square Regression (PLS-R)," *Vib. Spectrosc.*, vol. 116, p. 103288, 2021, doi: 10.1016/j.vibspec.2021.103288.
- [17] W. Tiaprasit and C. Sangpithukwong, *Buchi NIRFlex N-500 Training Course*. Bangkok: Buchi NIR Application Support, 2010.
- [18] H. Wang, X. Chu, P. Chen, J. Li, D. Liu, and Y. Xu, "Partial least squares regression residual extreme learning machine (PLSRR-ELM) calibration algorithm applied in fast determination of gasoline octane number with near-infrared spectroscopy," *Fuel*, vol. 309, p. 122224, 2021, doi: 10.1016/j.fuel.2021.122224.
- [19] M. M. Oliveira, J. P. Cruz-Tirado, J. V. Roque, R. F. Teófilo, and D. F. Barbin, "Portable near-infrared spectroscopy for rapid authentication of adulterated paprika powder," *J. Food Compos. Anal.*, vol. 87, p. 103403, 2020, doi: 10.1016/j.jfca.2019.103403.
- [20] D. Musaddad *et al.*, "The Potential of Chili Varieties as Powder Raw Material," *IOP Conf. Ser. Earth Environ. Sci.*, vol. 985, no. 1, 2022, doi: 10.1088/1755-1315/985/1/012044.
- [21] H. S. El-Mesery, H. Mao, and A. E. F. Abomohra, "Applications of non-destructive technologies for agricultural and food products quality inspection," *Sensors (Switzerland)*, vol. 19, no. 4, pp. 1–23, 2019, doi: 10.3390/s19040846.
- [22] B. T. Le, "Application of deep learning and near infrared spectroscopy in cereal analysis," *Vib. Spectrosc.*, vol. 106, p. 103009, 2020, doi: 10.1016/j.vibspec.2019.103009.
- [23] M. Zareef *et al.*, "An Overview on the Applications of Typical Non-linear Algorithms Coupled With NIR Spectroscopy in Food Analysis," *Food Eng. Rev.*, vol. 12, no. 2, pp. 173–190, 2020, doi: 10.1007/s12393-020-09210-7.
- [24] N. Okubo and Y. Kurata, "Non-destructive classification analysis of green coffee beans by using near-infrared spectroscopy," *Foods*, vol. 8, no. 2, 2019, doi: 10.3390/foods8020082.
- [25] G. B. Cagampang and F. M. Rodriguez, *Methods of Analysis For Screening Crops of Appropriate Qualities*. Los Banos, Philippines: Analytical Services Laboratory, Institute of Plant Breeding, University of The Philippines at Los Banos, 1980.
- [26] J. F. H. Jr, M. C. Howard, and C. Nitzl, "Assessing measurement model quality in PLS-SEM using confirmatory composite analysis," *J. Bus. Res.*, vol. 109, pp. 101–110, 2020, doi: 10.1016/j.jbusres.2019.11.069.
- [27] M. Sarstedt, C. M. Ringle, and J. F. Hair, *Partial Least Squares Structural Equation Modeling*. Switzerland: Springer International Publishing, 2021. doi: 10.1007/978-3-319-05542-8\_15-2.
- [28] F. D. Anggraeni, N. Khuriyati, M. A. F. Falah, H. Nishina, K. Takayama, and N. Takahashi, "Non-destructive measurement of lycopene content in high soluble solids stored tomato (*Solanum Lycopersicum* Mill. cv Rinka 409)," *Int. J. Adv. Sci. Eng. Inf. Technol.*, vol. 10, no. 6, pp. 256–2574, 2020, doi: 10.18517/ijaseit.10.6.9478.
- [29] R. Hayati, Z. Zulfahrizal, and A. A. Munawar, "Robust prediction performance of inner quality attributes in intact cocoa beans using near infrared spectroscopy and multivariate analysis," *Heliyon*, vol. 7, no. 2, p. e06286, 2021, doi: 10.1016/j.heliyon.2021.e06286.
- [30] R. N. M. J. Páscoa, P. A. L. S. Porto, A. L. Cerdeira, and J. A. Lopes, "The application of near infrared spectroscopy to wine analysis: An innovative approach using lyophilization to remove water bands interference," *Talanta*, vol. 214, p. 120852, 2020, doi: 10.1016/j.talanta.2020.120852.
- [31] M. A. Calegari, B. B. Ayres, L. M. dos Santos Tonial, S. M. de Alencar, and T. L. C. Oldoni, "Fourier transform near infrared spectroscopy as a tool for predicting antioxidant activity of propolis," *J. King Saud Univ. - Sci.*, vol. 32, no. 1, pp. 784–790, 2019, doi: 10.1016/j.jksus.2019.02.006.
- [32] O. O. Olarewaju *et al.*, "Model development for non-destructive determination of rind biochemical properties of 'Marsh' grapefruit using visible to near-infrared spectroscopy and chemometrics," *Spectrochim. Acta - Part A Mol. Biomol. Spectrosc.*, vol. 209, pp. 62–69, 2019, doi: 10.1016/j.saa.2018.10.027.
- [33] X. Jun-fang, L. Xiao-yu, L. Pei-wu, M. Qian, and D. Xiao-xia, "Application of Wavelet Transform in the Prediction of Navel Orange Vitamin C Content by Near-Infrared Spectroscopy," *Agric. Sci. China*, vol. 6, no. 9, pp. 1067–1073, 2007, doi: 10.1016/S1671-2927(07)60148-5.
- [34] P. Sodata and J. Peerapattana, "Application of near infrared spectroscopy with chemometrics for qualitative and quantitative dental caries assessment," *Vib. Spectrosc.*, vol. 111, p. 103170, 2020, doi: 10.1016/j.vibspec.2020.103170.
- [35] P. Sampaio, A. Soares, A. Castanho, A. S. Almeida, J. Oliveira, and C. Brites, "Dataset of Near-infrared spectroscopy measurement for amylose determination using PLS algorithms," *Data Br.*, vol. 15, pp. 389–396, 2017, doi: 10.1016/j.dib.2017.09.077.
- [36] L. R. Costa, G. H. D. Tonoli, F. R. Milagres, and P. R. G. Hein, "Artificial neural network and partial least square regressions for rapid estimation of cellulose pulp dryness based on near infrared spectroscopic data," *Carbohydr. Polym.*, vol. 224, p. 115186, 2019, doi: 10.1016/j.carbpol.2019.115186.
- [37] J. Jiang *et al.*, "Non-destructive quality assessment of chili peppers using near-infrared hyperspectral imaging combined with multivariate analysis," *Postharvest Biol. Technol.*, vol. 146, pp. 147–154, 2018, doi: 10.1016/j.postharvbio.2018.09.003.