

Characteristics of Anthocyanin as Natural Dyes from Butterfly Pea (*Clitoria ternatea* L.) on Regions Growing in North Maluku

Sri Soenarsih^a, Zauzah Abdullatif^a, Nurhasanah^b, Hamidin Rasulu^{c,*}

^a Department of AgroTechnology, Faculty of Agriculture, University of Khairun, Ternate, Indonesia

^b Departement Biology Education, Faculty of Teacher and Training Education, University of Khairun, Ternate, Indonesia

^c Department of Agricultural Product Technology, Faculty of Agriculture, Universitas Khairun, Ternate, Indonesia

Corresponding author: *hamidinrasulu@yahoo.com

Abstract—Butterfly pea has the potential to be a natural dye. The identification of anthocyanin compounds and anthocyanin content and antioxidant activity of butterfly pea flowers that grow in several regions in North Maluku need to be traced to identify the quality of anthocyanin content from late flowers. The research samples for the butterfly pea flower came from seven growing regions in North Maluku, and this study aimed to identify the phytochemical components of anthocyanins using the GC-MS method and to determine antioxidant activity using the DPPH method in ethanol extracts defined using the spectrophotometric method. The data obtained revealed the results of GC-MS phytochemicals and confectionery was identified as approximately 4-7 anthocyanin compounds with varying retention time peaks, with the best growth area for the highest anthocyanin content in Fitu Ternate with 35.38%, the Tidore region 33.92% and Maliaro 33.22%. Gas Chromatography-Mass Spectrometric (GC-MS) analysis results are intended to filter phytochemical, functional component groups presented by the Fourier Transform Infrared Spectrophotometer (FT-IR) in ethanol extract of *Clitoria ternatea* flowers. The results showed that the antioxidant activity test obtained IC₅₀ values on dried eagle flower extracts from several growing areas in North Maluku, which ranged from 39,009 – 74,481 µg / ml. IC₅₀ values are best produced in the growing area of Fitu Ternate and Tidore of 39,01 and 39,37 criteria for oxidant activity with powerful free radical inhibition.

Keywords— *Clitoria ternatea*; anthocyanins; natural dyes, antioxidants, growing regions.

Manuscript received 20 Aug. 2022; revised 21 May 2023; accepted 26 Jun. 2023. Date of publication 31 Aug. 2023. IJASEIT is licensed under a Creative Commons Attribution-Share Alike 4.0 International License.



I. INTRODUCTION

Among the countries that have plasma resources *nufah* high crop diversity is Indonesia. Many plants native to Indonesia are helpful and have not been excavated. *Clitoria ternatea*, or butterfly pea, is a plant native to Indonesia with a tropical climate. The first discovery of this flower was in the Ternate region, North Maluku Province, so the local people called it the local name blue flower because the flower is predominantly blue and functions as a natural dye. *Clitoria ternatea*, found in some growing areas in North Maluku, from this growing area, can provide the same or different anthocyanin content. The plant's growing area can also cause a high content of anthocyanins because agroclimatic and nutrient content in different soils can provide differences in anthocyanin content in flowers. *Clitoria ternatea* is a type of legume plant with purple or bluish-purple flowers. Blue or bluish-purple color on the flower can be used for coloring in food. Factory-made dyes (chemical) can be substituted with

dyes derived from plants obtained from *Clitoria ternatea*; the flower parts are blue or bluish-purple. These late petals produce anthocyanin color substances and are a group of flavonoid compounds used as natural dyes [1].

In *Clitoria ternatea*, there are components of anthocyanin compounds, so they have the potential as natural dyes in foodstuffs. The anthocyanin content of butterfly peas will be decisive in the color of the extraction results. Specifically, pelargonidin is a marker of orange to red, cyanidin indicates red, and delphinidin marks purple to blue. Likewise, Anthocyanin can function as a natural supplement, especially in boosting antibody (inflammatory), anti-microbial, anti-carcinogenic, and antioxidant, and includes anti-candida activity [2].

The color of the flower is blue; this color comes from anthocyanin compounds. The content of phytochemical compounds anthocyanins is stable and resistant to changes in acid-base colors that can be used as natural dyes for the culinary and food industries. The content of other phytochemicals in late flowers, such as flavonoids. The

flavonoid content of the flower can act as a source of warding off free radicals in the body because it acts as an antioxidant.

Given the huge potential of anthocyanins and the abundant content found in butterfly peas, *Antosian* is a compound derived from flavonoid glycosides, consisting of a sugar group (*glycon*), a non-glucose group, namely anthocyanidins (*glycocons*), and some anthocyanins containing acyl groups. But anthocyanins consist of *polyethoxy glycosylation* and *polyhydroxy* salt flavylum in their structure. Meanwhile, the anthocyanin form is in the form of *aglicon*, namely anthocyanins. However, the most widely distributed anthocyanins within the plant consist of delphinidin, pelargonidin, peonidin, petunidin, and cyaniding [3].

The potential of butterfly peas as a natural dye then the identification of anthocyanin compounds and anthocyanin content and antioxidant activity of butterfly peas that grow in several regions in North Maluku need to be traced to be identified the quality of anthocyanin content from butterfly pea flowers that grow in the North Maluku region.

II. MATERIAL AND METHOD

A. *Antosianin* Analysis

Samples of butterfly peas were taken from the regions of Bastiong Ternate, Tabona Ternate, Fitu Ternate, Maliaro Ternate, Kalaodi Tidore, Bibinoi Halmahera Selatan, Dehegila Morotai. The flowers are then cleaned of the dirt attached and continued with the process of drying flower samples using a cabinet drying at 40 °C. Then the sample that has been dried is mashed until it becomes a powder. The powder is sipped to produce a fine powder using 100 mesh for smooth and homogeneous powder. Solution maceration is done by making extracts of butterfly pea using the immersion method; the solution is then put into a *microtube*, then *vortexed* and *checkedrifuge*. The supernatant formed is used for GC-MS analysis methods. The analysis time is 60 minutes with an injector temperature of 260 °C, a detector of 250 °C, and a column of 325 °C. Furthermore, the flow rate using a 1 ml/min speed using a carrier media in helium gas. The identification process using the GC-MS method contained in the laboratory produces a list of compounds in which a mixture of gases and the molecular weight of each compound is presented as a chromatogram.

B. *Antioxidant* Analysis

Antioxidant activity with DPPH [4], the anthocyanins obtained by extracting are used to analyze DPPH compounds using a maceration technique ±24 hours, with the addition of a 96% ethanol solvent (1: 10), then centrifuged at 500 rpm, ± 14 minutes, the supernatant is obtained, and vacuum filtering is carried out. The filtrate is concentrated with an evaporator at 35 °C to obtain 1/10 of the initial filtrate as a concentrated extract. DPPH method can detect antioxidant activity and measurements using a UV-Vis spectrophotometer with λ 515 nm, where antioxidants are determined based on how much there is an absorption barrier in free radicals. Furthermore, calculate the percentage of inhibition of absorption of *1,1-diphenyl-2-picrilhydrazyl* (DPPH) using the formula:

$$\% \text{ of inhibition} = \frac{AO - A_{\text{sample}}}{AO} \times 100 \quad (1)$$

The percentage of inhibition obtained is a control absorbance (A0) or blank, while (A sample) is an absorbance of the test compound. In addition, 50% inhibition is a concentration of compounds that shows the calculation of the plot of the percentage of each inhibition to the concentration in the sample. For determining IC₅₀ (inhibition concentration), concentration variations range from 20% (b/v), 40% (b/v), 80% (b/v), 160% (b/v), Vitamin C as a positive control and extract that has the greatest percentage of inhibition at the initial treatment.

III. RESULT AND DISCUSSION

A. *Anthocyanin*

The anthocyanin content contained in the butterfly pea is characterized by a physical appearance in the form of blue pigments, and the results of previous studies have shown that this flower has benefits for human health. Anthocyanins in the pigment acid condition are reddish, while the pigment will show a blue hue in the alkaline condition. In addition, Anthocyanin is stable to high heat conditions and different pH, especially in coffee and acylation, which can maintain its structure well. The blue pigment of the flower is obtained by extracting the flower of the butterfly pea using a maceration method that produces a deep blue extract. Anthocyanins from flower flowers can be used as an acid-base indicator because these pigments change color when dripped with acids or bases and natural color substances.

Anthocyanin blue flower color will change color along with the change in pH value. At high pH, anthocyanins tend to be blue or colorless, then lead to red color at low pH. Most anthocyanins produce a color at a pH of less than 4. This discoloration is related to the hydroxy or methoxy group in the structure of the anthocyanin ring, thus affecting the color of anthocyanins. The dominant number of hydroxy groups causes the color to lead blue and is generally less stable. While the number of dominant methoxy groups compared to hydroxy groups in the structure of anthocyanins causes the color to be reddish.

Thus, in the GC-MS results, it can be seen that in the samples of butterfly peas from seven regions grown in North Maluku in ethanol extract, namely the *Bastiong*, *Tabona*, *Fitu*, *Maliaro*, *Tidore*, *Bacan*, and *Morotai* growing areas. From each area identified, approximately 4-7 anthocyanin compounds; the best growth area for the highest anthocyanin content is in Fitu Ternate with 35.38%, Tidore region with 33.92%, and Maliaro 33.22% and the rest in 4 growing regions with anthocyanin content ranging from 17.84% - 24.35%. The high content of anthocyanins in the three growing regions is likely because the region is a mountainous area, and the soil results from volcanic soil deposits with a lot of high enough nutrient content that can support the metabolism of anthocyanin formation in late flowers.

The study produced the number of monomeric anthocyanins expressed in mg equivalent *sianidin-3-glucoside* because Malvidin is one of the anthocyanins in the butterfly pea. The total content of Anthocyanin monomeric extracts from various regions grown in North Maluku consecutively is 21.55; 19.82; 35.38; 33.22; 33.92; 24.35; 17.84 (%) (bk). Total monomeric anthocyanins differed between samples of late confectionery from the area of origin

of growth studied. This difference is due to differences in the growing place and the environment. The biosynthesis process of anthocyanin compounds in each plant is strongly influenced by environmental conditions, especially air temperature, rainfall conditions, sunlight, and nutrients in the soil in each region where the plant grows [5].

It is reported that secondary metabolite compounds in phytochemical content, such as beta carotene and flavonoid components contained in plants, have different concentrations in which the plant grows. The components of such compounds are influenced by acidity (pH), environmental conditions, humidity, and nutritional elements contained in each soil nutrient [6]. The research report suggests cold environmental conditions are better suited for producing soybean anthocyanins and isoflavones [7]. According to [8], the higher the topographic plain where plants grow, the higher the anthocyanin content. [9] Reports of climate and growing places affect the virgin tread's bioactive components and antioxidant activity (*Catharanthus roseus*). It is also recommended that there are differences in the anthocyanin content of potatoes because they are influenced by the location where the plant grows. Still, the factor that has the most significant influence is the genotype factor of the potato plant on its anthocyanin content. It is also recommended that there are differences in the anthocyanin content of potatoes because they are influenced by the location where the plant grows. Still, the factor that has the most significant influence is the genotype factor of the potato plant on its anthocyanin content. [10], Moreover, the determining factors for the growth of plants with the best anthocyanin content are influenced by the biosynthesis process, mainly by the intensity of sunlight, good rainfall conditions, environmental conditions, and good nutrients in the soil or the area where the plant grows [5]. Thus, it can be concluded that the place of growth affects the monomeric anthocyanin production of butterfly peas. To better know the environmental influence on anthocyanin biosynthesis, further research is needed to pay close attention to the composition

of nutrients, the intensity of exposure to ultraviolet light, the temperature of day and night, as well as rainfall in the location and topography of the growth of late plants.

In addition, the stability of anthocyanins depends on light, pH, temperature, enzymes, metal ions, oxygen, anthocyanin pigments, antioxidants, and coffee. Anthocyanin compounds with good stability are dominated by a B ring contained in the anthocyanin content, and there is a *hydroxyl* or *methoxyl* group. Even lowering the stability of anthocyanins present in one solution has groups that can be known. Therefore, under different pH conditions, there will be a change in the color of several types of anthocyanin stability. Therefore, under different pH conditions, there will be a change in the color of various kinds of anthocyanin stability.

Furthermore, the pH condition of the sample greatly affects the anthocyanin color. Therefore, pH of the solution greatly affects the discoloration of the anthocyanin components. In addition, anthocyanins have a molecular structure with ionic properties [15]. Likewise, in acidic conditions, anthocyanin compounds appear to show a red color. Even anthocyanins show a purple hue with a neutral pH; on the contrary, under alkaline pH conditions, it turns blue. A large anthocyanin pigment consists of a red flavylium cation. Therefore, a lower pH of the solution will stabilize the anthocyanin compounds. In addition, flavylium cations with a low pH will affect anthocyanin compounds and be water-soluble. A decrease in water concentration increases the rate of deprotonation of flavylium cations, thereby lowering the stability of the color. In addition to pH, the polymerization of anthocyanins can also improve color stability at lower pH.

The extraction results using *ethanol* on *Clitoria ternate* using GC-MS analysis will obtain the appearance of phytochemical component profiles. The results of FT-IR analysis also show functional groups with their grouping. Ethanol extract with GC-MS reveals that *Clitoria ternate* has many components of the active compound (Table 1 and Figure 1).

TABLE I
THE COMPOUND WAS IDENTIFIED BY GC-MS ANALYSIS USING ETHANOL EXTRACT *C. TERNATEA* FROM NORTH MALUKU

Areas Sample	Retention Times	Name of the Compounds	Formula Molecular	Molecular weight (kDa)	Area Peak	Anthocyanin Content (%)
Bastiong	12,241	Melezitose	C ₁₈ H ₃₂ O ₁₆	504	0.38	21,55
	15,513	Melezitose	C ₁₈ H ₃₂ O ₁₆	504	0.37	
	18,087	Melezitose	C ₁₈ H ₃₂ O ₁₆	504	11.15	
	21,604	Melezitose	C ₁₈ H ₃₂ O ₁₆	504	5.84	
	29,668	L- (+)- ascorbic acid 2,6-dihexadecanoate	C ₃₈ H ₆₈ O ₈	652	3.57	
	32,260	9-octadecenoic acid (Z)-2-hydroxyl-1(hydroxymethyl) ethl ester	C ₂₁ H ₄₀ O ₄	356	0.24	
Tabona	15,512	Melezitose	C ₁₈ H ₃₂ O ₁₆	504	0.42	19,82
	17,478	Eicosanoic acid	C ₂₀ H ₄₀ O ₂	312	0.51	
	18,009	Melezitose	C ₁₈ H ₃₂ O ₁₆	504	10.93	
	22,318	Melezitose	C ₁₈ H ₃₂ O ₁₆	504	0.96	
	29,671	L- (+)- ascorbic acid 2,6-dihexadecanoate	C ₃₈ H ₆₈ O ₈	652	7.00	
	10,060	Melezitose	C ₁₈ H ₃₂ O ₁₆	504	1.11	
Fitu	15,509	Melezitose	C ₁₈ H ₃₂ O ₁₆	504	0.36	35.38
	18,022	Melezitose	C ₁₈ H ₃₂ O ₁₆	504	13.29	
	22,325	Melezitose	C ₁₈ H ₃₂ O ₁₆	504	2.31	
	29,671	Melezitose	C ₃₈ H ₆₈ O ₈	652	6.29	

Areas Sample	Retention Times	Name of the Compounds	Formula Molecular	Molecular weight (kDa)	Area Peak	Anthocyanin Content (%)	
Maliaro		L- (+)- ascorbit acid 2,6-dihexadecanoatee					
	32,980	Oleic acid	C ₁₈ H ₃₄ O ₂	282	12.02		
	12,237	1- Decanol, 2- ethyl Melezitose	C ₁₂ H ₂₆ O	186	0.58		
	15,516	Melezitose	C ₁₈ H ₃₂ O ₁₆	504	0.55		
	16,125	Melezitose	C ₁₈ H ₃₂ O ₁₆	504	0.21		
	17,505	Melezitose	C ₁₈ H ₃₂ O ₁₆	504	0.65	33.22	
	18,104	L- (+)- ascorbic acid 2,6-dihexadecanoate	C ₁₈ H ₃₂ O ₁₆	504	13.46		
	29,678	Oleic acid	C ₃₈ H ₆₈ O ₈	652	8.64		
	Bacan	32,980		C ₁₈ H ₃₄ O ₂	282	9.13	
		15,516	Melezitose	C ₁₈ H ₃₂ O ₁₆	504	0.49	
18,104		Melezitose	C ₁₈ H ₃₂ O ₁₆	504	13.32		
19,655		Melezitose	C ₁₈ H ₃₂ O ₁₆	504	0.65		
22,372		Melezitose	C ₁₈ H ₃₂ O ₁₆	504	0.92		
29,678		L- (+)- ascorbic acid 2,6-dihexadecanoat	C ₃₈ H ₆₈ O ₈	652	8.55	24.35	
39,221		9-octadecenoic acid (Z)-2-hydroxyl-1(hidroxyethyl) ester	C ₂₁ H ₄₀ O ₄	356	0.42		
Morotai	1,816	Eicosanoic acid	C ₁₈ H ₃₂ O ₁₆	504	0.14		
	15,509	Melezitose	C ₁₈ H ₃₂ O ₁₆	504	0.28		
	17,961	Melezitose	C ₁₈ H ₃₂ O ₁₆	504	6.67		
	21,876	Melezitose	C ₁₈ H ₃₂ O ₁₆	504	1.28		
	29,678	L- (+)- ascorbic acid 2,6-dihexadecanoat	C ₃₈ H ₆₈ O ₈	652	9.17	17.84	
	32,283	9-octadecenoic acid (Z)-2-hydroxyl-1(hidroxyethyl) ester	C ₂₁ H ₄₀ O ₄	356	0.30		




Fig. 1 GC-MS chromatogram of anthocyanin compounds present from ethanol extract *Clitoria ternatea*

B. Antioxidant Activity

Bars Research results to find out the results of extraction of dried flowers using ethanol solvent; the use of the DPPH method in analyzing antioxidant activity based on the ability to use ethanol extract and can reduce free radical compounds, especially DPPH compounds. This study used samples of butterfly peas taken from growing regions in four districts in North Maluku. In the extraction process, the flower sample is cleaned of dirt, dried, cut into small pieces, and put into a

bottle containing 96% ethanol solvent. The solvents' solubility and the dissolved components are the basis of adding solvents to a material. In this study, ethanol solvents were used. This solvent was chosen because, according to [12], [13], ethanol can extract more active compounds than other types of organic solvents, besides the utilization of this confectionery is intended for natural dyes for food, so it is expected not to cause toxins that can cause poisoning when used for food and relatively safer.

TABLE II
CLITORIA TERNATEA FLOWER EXTRACT ON ANTIOXIDANT ACTIVITY FROM NORTH MALUKU

Region	IC ₅₀ (µg/ml)		Average IC ₅₀ (µg/ml) ±SD	Antioxidant Category
	Rep 1	Rep 2		
Bastiong	60.86	62.83	61.84 ± 1.39	Strong
Tabona	73.68	75.29	74.48 ± 1.14	Strong
Fitu	39.62	39.11	39.37 ± 0.36	Very Powerful
Maliaro	59.07	53.98	56.51 ± 3.62	Strong
Tidore	38.91	39.11	39.01 ± 0.15	Very Powerful
Bacan	63.47	52.46	57.96 ± 7.79	Strong
Morotai	59.00	59.58	59.29 ± 0.40	Strong

Based on data from the analysis of antioxidant activity, IC₅₀ values were obtained in dried confectionery extracts from several growing regions in North Maluku, which ranged from 39,01 to 74,48 µg/ml. IC₅₀ values are best produced in the growing area of Fitu Ternate and Tidore of 39.01, and 39.37 criteria for oxidant activity are very strong. This shows that samples of butterfly peas that grow in the North Maluku region have a strong to very strong level of antioxidant activity. The concentration contained in the sample solution has an IC₅₀ value by reducing DPPH activity by 50%, where the smaller the IC₅₀ value, the higher the reading of antioxidant activity in the sample [14], [15], and the more antioxidant activity to reduce the stronger. Table 3 presents antioxidant levels using the DPPH method at IC₅₀ values.

TABLE III
LEVELS OF ANTIOXIDANT POWER WITH THE DPPH METHOD

Value IC ₅₀ (ppm)	Level of Antioxidant Activity
151-200	Weak
100-150	Is
50-100	Strong
<50	Very Strong

Source [31]

In the test results using dry *Clitoria ternatea* extract obtained, the results of the antioxidant activity test in Table 3 show that the susceptibility of IC₅₀ values < 100 ppm is a strong and very strong activity. This is due to the analysis of the low water content of the butterfly pea sample. Water content is low enough when analyzing the activity of the ingredients causing antioxidant compounds contained in the ingredients will be higher in value the water content is low, then the content of antioxidant compounds can be detected better, meaning that antioxidant activity will be stronger if the water content of the ingredient is reduced or low [16], [23]-[24]. When the water content of the material is high, it can cause flavonoid compounds to oxidize, thereby decreasing antioxidant activity.

Using the DPPH method in butterfly pea extract will show results where the antioxidant activity is very strong, likely due to the presence of phenolic compounds in it. The antioxidant mechanism of phenolic compounds is based on oxidation-reduction reactions. Due to reducing agents contained in this phenolic compound, there is a reduction in reactive free radical compounds and reactions between DPPH radicals and phenolic compounds in late flowers as radical capture compounds [25], [26].

Meanwhile, to interpret the results of the DPPH method, it was recently introduced, namely efficient concentration (EC₅₀), or called the IC₅₀ value, as one of the parameters in calculating antioxidants. IC₅₀ can be interpreted as a substrate

concentration that can eliminate 50% of DPPH activity through discoloration. Therefore, several research results show that phenolic compounds strongly correlate with antioxidant activity [17], [18], [27], [28].

Phenolics in ethanol extract will release H•, one of the free radicals. H• will bind to the DPPH radical to form a new compound, the stable *diphenyl picrilhydrazin*. Phenolic compounds contained in ethanol extract as a free radical catcher that loses H will become new radicals that are relatively more stable and harmless to the body because of the resonance effect of antiaromatic so that free radicals are not formed and can prevent or repair tissue damage that is the effect of free radical attacks. Phenolic compounds from heterogeneous chemical groups containing phenol groups (functional hydroxyl groups in aromatic rings) in their basic structure. In addition, phenolic compounds can induce antioxidant protein synthesis and increase the activity of antioxidant enzymes [19], [20].

Antioxidant activity in the butterfly pea (*Clitoria ternatea*) calculation of IC₅₀ value with linear regression curve, so that a concentration with the percentage of damping in the test sample is obtained and then made in a linear regression curve, $y = bx + a$, where x is the concentration (ppm), and y is the percentage of IC₅₀, where the relationship between the concentration of the test solution with percent damping, obtained regression equation with $R^2 = 0.8216$ from the Bastiong Ternate region. From the value of R^2 , it can be known that there is a significant relationship between the concentration of the solvent and the percentage of immersion observed with a degree of intensity of 0.8216; this indicates an 82% degree of inhibition (Figure 2). The highest inhibition power in the growing area Fitu Ternate amounted to $R^2 = 0.9525$ degrees inhibition by 0.95%, and the growing area Tidore $R^2 = 0.9499$ degrees inhibition 94% is powerful. Other growing regions obtained R2 values ranging from 0.7877 - 0.8343 with degrees of inhibition of 78 % - 83% criteria for strong inhibitory antioxidant activity.

These results align with previous researchers' findings that found that the reducing power of anthocyanins of purple sweet potato extract [21], [22], and lychee skin anthocyanin was greater than that of ascorbic acid [29]. For this reason, it can be concluded that the growing condition of *Clitoria Ternatea* does not affect the anthocyanin content in the form of antioxidant activity. The degree of similarity in carrying out anthocyanin extraction affects the antioxidant activity produced. Although the reducing force and free radical capture activity are at pH 1 because it is the more dominant *flavilium cation* compound in the anthocyanin structure, it can

provide hydrogen cations that easily reduce Fe^{3+} to Fe^{2+} and stabilize DPPH electrons [30].




Fig. 2 Ethanol Extract of *Clitoria Ternatea* with Antioxidant Activity from Ternate Bastiong




Fig. 3 Ethanol Extract of *Clitoria Ternatea* with Antioxidant Activity from Ternate Tabona




Fig. 4 Ethanol Extract of *Clitoria Ternatea* with Antioxidant Activity from Ternate Fitu




Fig. 5 Ethanol Extract of *Clitoria Ternatea* with Antioxidant Activity from Ternate Maliaro




Fig. 6 Ethanol Extract of *Clitoria Ternatea* with Antioxidant Activity from Tidore




Fig. 7 Ethanol Extract of *Clitoria Ternatea* with Antioxidant Activity from Bacan




Fig. 8 Ethanol Extract of *Clitoria Ternatea* with Antioxidant Activity from Morotai

IV. CONCLUSION

Analysis of GC-MS phytochemicals was identified as approximately 4-7 anthocyanin compounds; the best growth area for the highest anthocyanin content is in Fitu Ternate with 35.38%, Tidore region 33.92% and Maliaro 33.22%. Therefore, in this study, the results of testing antioxidant activity obtained IC_{50} values from dried butterfly pea extract from several growing areas in North Maluku, which ranged from 39,01 – 74,48 µg/ml. IC_{50} values are best produced in the growing area of Fitu Ternate and Tidore of 39,01 and 39,37 criteria for oxidant activity with very strong free radical inhibition.

ACKNOWLEDGEMENT

This research is fully supported by the Funding of Competitive Research Excellence of Universities (PKUPT) Khairun University in 2021 No:016 / PEN-PKUPT / PG.12 / 2021.

REFERENCES

- [1] H. Hariadi, M. Sunyoto. B. Nurhadi and A. Karuniawan. "Comparison of phytochemical characteristics pigmen extract (Antosianin) sweet purple potatoes powder (*Ipomoea batatas* L) and clitoria flower (*Clitoria ternatea*) as natural dye powder". *Journal of Pharmacognosy and Phytochemistry*. 7(4): 3420-3429. 2018.
- [2] A.Q. Syafa'atullah, A. Arie Amira, S. Sonya Hidayati and M. Mahfud. "Anthocyanin from butterfly pea flowers (*Clitoria ternatea*) by ultrasonic-assisted Extraction". *AIP Conference Proceedings*. 2020.
- [3] B. Anthika, S.P. Kusumocahyo, and H. Sutanto. ""Ultrasonic approach in Clitoria ternatea (butterfly pea) extraction in water and extract sterilization by ultrafiltration for eye drop active ingredient". *Procedia Chemistry*, 16, 237-244. 2015.
- [4] N. M. Thuy, V. Q. Minh, T. C. Ben, M.T. Thi Nguyen, H.T. N. Ha, and N. V. Tai. "Identification of anthocyanin compounds in butterfly pea flowers (*C. ternatea* L.) by ultra performance liquid chromatography/ultraviolet coupled to mass spectrometry". *Molecules*, 26(15), 4539. 2021.
- [5] P. Kumari and B. Bhargava. "Phytochemicals from edible flowers: Opening a new arena for healthy lifestyle". *Journal of Functional Foods*, 78, 104375. 2021.
- [6] H. E. Khoo, A. Azlan, S.T. Tang, and S.M. Lim. "Anthocyanidins and anthocyanins: Coloured pigments as food, pharmaceutical ingredients, and the potential health benefits". *Food & nutrition research*, 61(1), 1361779. 2017.
- [7] E. H. Kim, O.K. Lee, J.K. Kim, S.L. Kim, J. Lee, S.H. Kim, I.M.M. Chung. "Isoflavones and anthocyanins analysis in soybean (*Glycine max* (L.) Merrill) from three different planting locations in Korea". *Field Crops Res* 156: 76–83. 2014.
- [8] J. H. Y. Galani, P. M. Mankad, A. K. Shah, N. J. Patel, R. R Acharya, and J.G. Talati, "Effect of storage temperature on vitamin C, total phenolics, UPLC phenolic acid profile and antioxidant capacity of eleven potato (*Solanum tuberosum*) varieties". *Horticultural Plant Journal*, 3(2), 73-89. 2017.
- [9] A. Kumar, K.C. Singhal, R.A. Sharma, K.V. Govind and V. Kumar. "Total Phenolic and Antioxidant Activity of in Different Geographical Locations of Rajasthan Catharanthus Roseus". *Asian J. Exp. Biol. Sci.* Vol 4 (1): 155-158. 2014.
- [10] G. C. V. Gamage, Y. Y. Lim, and W. S.. Choo. "Anthocyanins from Clitoria ternatea flower: Biosynthesis, extraction, stability, antioxidant activity, and applications". *Frontiers in Plant Science*, 12. 2021.
- [11] T. N. Pham, X. T. Le, P. T. N. Nguyen, T. H. Tran, T. P. Dao, D. H. Nguyen, and H. L. T. Anh, "Effects of storage conditions on total anthocyanin content of Butterfly pea flower (*Clitoria ternatea* L.)". In *IOP Conference Series: Materials Science and Engineering* (Vol. 736, No. 6, p. 062005). IOP Publishing. 2020.
- [12] T. T. Vuong and P. Hongsrabhas. "Influences of pH on binding mechanisms of anthocyanins from butterfly pea flower (*Clitoria ternatea*) with whey powder and whey protein isolate". *Cogent Food & Agriculture*, 7(1), 1889098. 2021.
- [13] S. Lin, X. Meng, C. Tan, Y. Tong, M.Wan, M.Wang and Y. Ma. "Composition and antioxidant activity of anthocyanins from *Aronia melanocarpa* extracted using an Ultrasonic-Microwave-Assisted Natural Deep Eutectic Solvent extraction method". *Ultrasonics Sonochemistry*, 106102. 2022.
- [14] J. S. de Moraes, A. S. Sant'Ana, A. M. Dantas, B. S. Silva, M.S. Lima, G.C. Borges and M. Magnani, "Antioxidant activity and bioaccessibility of phenolic compounds in white, red, blue, purple, yellow and orange edible flowers through a simulated intestinal barrier". *Food Research International*, 131, 109046. 2020.
- [15] Z. Sapiun, P. S. Wicita, A. S. Buloto, V. Kamba, A.K. Imran, W. A. A. Buana and Y. Banne. "Preparation and Characterization of Butterfly Pea (*Clitoria ternatea* L.) Nanoemulgel". *Open Access Macedonian Journal of Medical Sciences*, 10(A), 585-589. 2022.
- [16] A. E. Al-Snafi, "Pharmacological importance of *Clitoria ternatea*—A review". *IOSR Journal of Pharmacy*, 6(3), 68-83. 2016.
- [17] E. J. Jeyaraj, Y. Y. Lim and W. S. Choo. "Extraction methods of butterfly pea (*Clitoria ternatea* L) flower and biological activities of its phytochemicals". *Journal of food science and technology*, 58(6), 2054-2067. 2021.
- [18] D.Y. Zhang, Yi wan, Run-Ze Hu, Jing-Hi Hao. "Evaluation of the alkaloid, polyphenols, and antioxidant contents of various mulberry cultivars from different planting areas in eastern China". *Industrial Crops and Products*, 15 out. 2018. v.122, p.298–307. 2018.
- [19] B. Srichaikul, "Ultrasonication extraction, bioactivity, antioxidant activity, total flavonoid, total phenolic and antioxidant of *Clitoria ternatea* linn flower extract for anti-aging drinks". *Pharmacognosy Magazine*, 14(56), 322. 2018.
- [20] G. D'Urso, J. M. Jurriaan P. Paola Montoro, D.H. Robert and C.H. Ric de Vos. "Identification of Bioactive htyochemicals in Mulberries". *Metabolites* Vol 10 (7): 2-16. 2020.
- [21] P. de Aguiar Cipriano, H. Kim, C. Fang, V. P. Venancio, S.U. Mertens-Talcott and S. T. Talcott. "In vitro digestion, absorption and biological activities of acylated anthocyanins from purple sweet potatoes (*Ipomoea batatas*)". *Food Chemistry*, 374, 131076. 2022.
- [22] S. A. T. Lakshan, N. Y. Jayanath, W. P. K. M. Abeysekera and W. K. S. M. Abeysekera, "A commercial potential blue pea (*Clitoria ternatea* L.) flower extract incorporated beverage having functional properties". *Evidence-Based Complementary and Alternative Medicine*, 2019.
- [23] X. Si, Q. Chen, J. Bi, X. Wu, J. Yi, L. Zhou and Z. Li. "Comparison of different drying methods on the physical properties, bioactive compounds and antioxidant activity of raspberry powders". *Journal of the Science of Food and Agriculture*, 96(6), 2055-2062. 2016.
- [24] M. Al-Duais, L. Müller, V. Böhm and G. Jetschke, "Antioxidant capacity and total phenolics of *Cyphostemma digitatum* before and after processing: use of different assays". *European food research and technology*, 228(5), 813-821. 2009.
- [25] M. G. Morais, A. A. Saldanha, L. S. Azevedo, I. C. Mendes, J. P. C. Rodrigues, P. A. Amado and L. A. R. dos Santos Lima. "Antioxidant and anti-inflammatory effects of fractions from ripe fruits of *Solanum lycocarpum* St. Hil.(Solanaceae) and putative identification of bioactive compounds by GC–MS and LC-DAD-MS". *Food Research International*, 156, 111145. 2022.
- [26] N. Jiménez, L. Carrillo-Hormaza, A. Pujol, F. Álzate, E. Osorio and O. Lara-Guzman. "Antioxidant capacity and phenolic content of commonly used anti-inflammatory medicinal plants in Colombia". *Industrial Crops and Products*, 70, 272-279. 2015.
- [27] Z. Shang, M. Li, W. Zhang, S. Cai, X. Hu and J. Yi. "Analysis of phenolic compounds in pickled chayote and their effects on antioxidant activities and cell protection". *Food Research International*, 157, 111325. 2022.
- [28] C. Cirak, J. Radusiene, L. Raudone, G. Vilkičkyte, F. Seyis, M. Marksas, and F. Yayla. "Phenolic compounds and antioxidant activity of *Achillea arabica* populations". *South African Journal of Botany*, 147, 425-433. 2022.
- [29] G. Toydemir, B. G. Subasi, R. D. Hall, J. Beekwilder, D. Boyacioglu, and E. Capanoglu. "Effect of food processing on antioxidants, their bioavailability and potential relevance to human health". *Food Chemistry: X*, 100334. 2022.
- [30] H. M. Ali, W. Almagribi and M. N. Al-Rashidi. "Antiradical and reductant activities of anthocyanidins and anthocyanins, structure–activity relationship and synthesis". *Food Chemistry*, 194, 1275-1282. 2016.
- [31] Akbari, A., Izadi-Darbandi, A., Bahmani, K., Farhadpour, M., Ebrahimi, M., Ramshini, H., & Esmaeili, Z. Assessment of phenolic profile, and antioxidant activity in developed breeding populations of fennel (*Foeniculum vulgare* Mill). *Biocatalysis and Agricultural Biotechnology*, 48, 102639. 2023