

The swab specimens were aseptically placed into 3 milliliters of viral transport medium, namely the CITOSWAB VTM collection and transportation kit. In July in the year 2021, the esteemed Central Laboratory of Universitas Padjajaran conducted the preparation of samples and extraction of RNA utilizing a biosafety level-3 (BSL-3) laboratory. The viral RNA was obtained using a column-based RNA/DNA extraction kit (catalog number SPM-48, Da-An Gene Co., Ltd.). Subsequently, we employed a real-time reverse transcription polymerase chain reaction (RT-PCR) approach that specifically targeted the open reading frame 1ab (ORF 1ab) and nucleocapsid (N) genes, utilizing the PCR Fluorescence Probing technique (Catalog number DNC-96, Da-An Gene Co., Ltd), under the manufacturer's prescribed protocols, to assess the integrity of the SARS CoV-2 specimen. The UJI COVID-19 IgG/IgM kit (Catalogue Number AUC-10D20, manufactured by Pakar Biomedika Indonesia) was employed to assess the efficacy of the SARS-CoV-2 antibody.

III. RESULTS AND DISCUSSIONS

A. Results

Upon careful examination of the data presented in Table 1, it has come to our attention that among the cohort of 19 felines under investigation, a total of four individuals (namely, Cat7, Cat10, Cat12, and Cat13) have yielded positive results for SARS-CoV-2 upon conducting nasal, oropharynx, and rectum swabs, as illustrated in Figure 1, Figure 2, and Table 1.

TABLE I
THE NASAL, OROPHARYNX AND RECTUM SWAB OF SUSPECTED FELINES
CHARACTERISTICS OF FELINES WITH POSITIVE COVID-19 OWNERS IN
BANDUNG, INDONESIA

Felines	Swab		
	Nasal	Oropharynx	Rectum
1	-	-	-
2	-	-	-
3	-	-	-
4	-	-	-
5	-	-	-
6	-	-	-
7	+	+	+
8	-	-	-
9	-	-	-
10	+	+	+
11	-	-	-
12	+	+	+
13	+	+	+
14	-	-	-
15	-	-	-
16	-	-	-
17	-	-	-
18	-	-	-
19	-	-	-

Notably, the incidence rate observed in the present study surpasses that of the study conducted in Spain, wherein SARS-CoV-2 was detected in 12.5% of the felids scrutinized. Two felids that underwent SARS-CoV-2 testing (Cat7 being asymptomatic, while Cat10, 12, and 13 exhibiting flu-like symptoms) had intimate associations with their respective owners, involving physical proximity and cohabitation. The

felids can potentially contract SARS-CoV-2 from asymptomatic carriers [23]. Notably, out of the 38 RT-qPCR swab data, twelve were deemed inconclusive, resulting in a percentage of 31%. It is plausible that the inconclusive outcomes could be attributed to the sole detection of either the N-Gene or Orf1ab. The N-gene, situated at the apex of the genetic hierarchy, confers protection to the enclosed genetic material by ensconcing it within a protective shell. Acknowledging that the N-Gene plays a crucial role in safeguarding the viral RNA gene center in the context of SARS-CoV-2 is imperative. It is imperative to highlight that the predominant gene of this virus is Orf1ab, which exhibits open reading frames with overlapping characteristics [24,25]. The indeterminate outcomes may have arisen due to SARS-CoV-2 identification fragments, contaminants, or a procedural aberration during the initial phase of sample handling. Given that we have conducted RT-qPCR on two separate occasions and obtained concordant outcomes, it appears plausible that there were no technical aberrations during the RT-qPCR procedure.




Fig. 2 Diagnostic test results from a suspected cat sample




Fig. 3 Positive results marked by 2 lines appearance of the kit, indicate the presence of viral antigens.

Regrettably, the serological assay failed to yield any affirmative outcomes. The findings suggest that the rapid antibody detection kit could not identify the occurrence of SARS-CoV-2 antibody, and that felids may have exhibited no antibody-antigen response to SARS-CoV-2 or that the antibody had already undergone clearance from their bloodstream. The findings of a prior investigation have demonstrated that felids possess neutralizing antibodies against SARS-CoV-2, with titers ranging from 1/20 to 1/1080. In order to elucidate the intricacies of the antibody-

antigen interplay of SARS-CoV-2, particularly in the context of domesticated fauna, further experimentation is imperative.

B. Discussion

The risks for zoonotic animals to humans' transmission of COVID-19, particularly in the context of companion animals such as canines, felines, and other domesticated species, underscores the need for heightened attention to this matter. This issue calls for further examination and reflection within the domain of epidemiology. It has been duly noted that the SARS-CoV Type 2 virus exhibits a significant affinity towards domesticated *Canidae* and *felidae* species that cohabit with at least one individual diagnosed with the virus, as per the latest research in the field of environmental health. As mentioned earlier, the statement postulates the risks that the disease could be passed from animals to humans or vice versa [27]. Scholars have postulated that the diverse suppositions regarding the genesis of SARS-CoV-2 are presently unverifiable or falsifiable. Upon conducting a thorough phylogenetic analysis, it has been ascertained that SARS-CoV Type 2 exhibits a significantly heightened degree of genetic resemblance, amounting to 96%, with two SARS-like coronaviruses, namely bat SL CoVZX45 and bat SL CoVZX21.

Notably, both of these coronaviruses are naturally harbored by chiropteran species, which underscores the importance of understanding the complex interactions between wildlife and human health. The SARS-CoV Type 2 pathogen exhibits a proclivity for binding to particular receptors, thereby facilitating its invasion of horseshoe bats and civets. It has been duly noted that the susceptibility of gallinaceous avifauna, porcine taxa, anatine avifauna, and murine rodents to this particular pathogen appears to be inconsequential [23]. It is widely acknowledged within the scientific community that the Feline Coronavirus (FCoV) group acts as the precursor virus of the Feline Infectious Peritonitis Virus (FIPV), a pathogen that is highly prevalent among domesticated felines worldwide [31]. On the contrary, a proposal has been posited positing that the origin of group II Feline Coronavirus (FCoV) may be ascribed to the amalgamation of canine and feline CoVs. From an epidemiological perspective, it appears that this hypothesis is relatively less plausible and demonstrates a significant biological and molecular pathogenesis compared to cases involving Feline Coronavirus type I. As a senior professor in environmental health, it is important to consider all possible factors and analyze the evidence thoroughly before drawing any conclusions.

The results of a recent investigation suggest that the contagion of the SARS-CoV Type 2 virus in felines is not of significant concern. Regarding felines, it is noteworthy that the transmission of the disease exhibits a remarkable degree of resilience, thereby facilitating contagion to susceptible felines through droplet dissemination. It has been duly noted that a considerable proportion of animal species that are susceptible to SARS-CoV Type 2 exhibit modifications in various amino acids of their ACE2 proteins, as per the current scientific literature. It is hypothesized that these specific taxonomic groupings, particularly those in close proximity to human habitation, demonstrate an increased vulnerability to contracting the SARS-CoV Type 2 pathogen. It is conceivable

to postulate that in the event of a robust establishment by one of these organisms, it may potentially act as a reservoir for the dissemination of disease among other wildlife, as has been previously proposed in scholarly literature [34]. After conducting sequencing research on ACE2 from several mammals, it has been observed that the domestic cat (*Felis catus*) displays the highest degree of sequence similarity, with an overall identity of 85.2%. It is noteworthy to observe that the *Manis javanica*, commonly known as the Malayan pangolin, and *Oryctolagus cuniculus*, which is referred to as the European rabbit, display a remarkably analogous genomic constitution, with a similarity rate of 84.8% between the two taxa. The comparative analysis of these findings has been conducted under established protocols and methodologies in the field of zoonoses research, utilizing the human ACE2 sequence.

The latest scientific findings have demonstrated that the SARS-CoV Type 2 pathogen ACE2 as a receptor for cellular entrance, as has been evidenced. It has been duly noted that the felid species demonstrate a comparatively elevated expression of ACE2, particularly in their tissues, digestive tract, and urinary system, in contrast to the human populace. Notably, felids demonstrate a greater inclination towards acquiring and transmitting illnesses through various means, in contrast to canines [37].

It has been postulated with significant frequency that domestic felids represent the domesticated species that are most commonly affected by SARS-CoV Type 2 infection. The findings of the RT-PCR analysis reveal the detection of SARS-CoV-2 in six felines that have been procured from various geographical regions, including France, Spain, Chile, Hong Kong, and New York. Within the cohort of six felines presently under investigation, a noteworthy proportion of four individuals have displayed clinical manifestations, including emesis, diarrhea, dyspnea, asthenia, anorexia, cough, sneezing, and ocular discharge. Following the confirmation of the individual's positive status for Covid 19, clinical manifestations were observed on the seventeenth day of the disease course.

There have been several documented instances of COVID-19 infections in both human and non-human animal populations, including felids. It has been postulated that felids may have acquired the infection from their human caregivers who were asymptomatic at the time, as per the literature citation 23. It is anticipated that future research endeavors will augment our understanding of the transmission dynamics of the disease, particularly in the realm of zoonotic transmission. It is of utmost importance to provide additional clarification in order to develop and execute effective preventative measures aimed at reducing the transmission of COVID-19 [35].

Regarding transmission, it is essential to recognize that the SARS-CoV-2 virus employs a receptor similar to that of the respiratory mucosa. This phenomenon increases the likelihood of zoonotic virus transmission [36]. A recent investigation suggests that SARS-CoV-2-infected felids are capable of passing the virus to other felids through direct contact. Nevertheless, the extent of their potential to pass the virus to other animals or humans remains unclear, necessitating additional research [40]. Due to the homology of nucleotides and amino acids within the spike protein, it is

plausible that inter-animal transmission could occur despite the uncertainty surrounding the concept of zoonotic virus transmission. Notable similarity exists between the genomes of SARS-CoV Type 2 and other animal coronaviruses. The nucleotides may constitute the genetic material changed during transmission to animal hosts. It is plausible that variations in the amino acid sequences may have resulted, possibly imparting increased virulence to certain taxa, particularly those containing the spike protein.

According to the evidence, the likelihood of zoonotic transmission appears relatively low. Given the remarkable similarity between the nucleotides and amino acids of the spike protein of SARS-CoV Type 2 and those of other animal coronaviruses, it is plausible that the virus could be transmitted between species. It is possible that nucleotide mutations occurred during zoonotic transmission of the virus to animal hosts expressing spike protein-associated amino acids, thereby increasing the pathogenicity of the virus in said animal hosts [10].

In the context of SARS-CoV and SARS-CoV-2 infection in cats, it is notable that the clinical manifestations are typically limited to moderate respiratory symptoms. It has been observed that felids infected with SARS CoV-2 exhibit viral RNA in their pulmonary organs and gastrointestinal tracts for up to 6 days and 3 days, respectively, following infection. In addition, it is noteworthy that on the third post-infection day, viral ribonucleic acid was detected only in the pulmonary tissue of these felids. The relevant information is documented in reference 35. However, there have been documented instances of viral transmission via the fecal-oral route [41]. The pathogen SARS CoV-2 primarily causes infections that are restricted to the upper respiratory tract. The general public has hypothesized that feline infections can be attributed to their human caregivers. However, no empirical evidence has established the plausibility of interspecies or zoonotic transmission of infections. [35]

According to previous research, the present findings indicate that felids that have been exposed to sick individuals in the past may have detectable levels of SARS-CoV-2 [22]. According to the findings of reference 41, SARS CoV-2 has the propensity to replicate efficiently in feline species and can be transmitted between felines via droplets. Given the ambiguity surrounding the transmission dynamics of the virus between humans and felines, or vice versa, it is imperative to emphasize that the positive results cannot be interpreted as conclusive evidence of feline susceptibility to SARS-CoV-2. If the answer is affirmative, the potential transmission of SARS-CoV-2 from human caregivers to their feline counterparts should be considered as an alternative scenario. It is highly recommended to conduct additional in vitro analyses using gastrointestinal tract-derived cell lines to address this issue.

During the course of our investigation, we encountered a number of obstacles. The limitations of our investigation included several factors, such as the sample size, which, despite being adequate, could have been larger. In addition, the number of individuals who owned SARS-CoV-2-infected cats was limited, limiting the generalizability of our findings. The timing of specimen collection was also problematic, as it affected both PCR and serologic results. In addition, the lack of a feline-specific COVID-19 IgG/IgM test was a significant

limitation we faced. The imposition of a curfew in the study area restricted the mobility of our personnel, which constituted an additional difficulty. This has substantially hindered our collaborative efforts to investigate potential human-animal transmission routes.

Our findings indicate that physical contact or cohabitation between an infected cat owner and another feline companion may increase the likelihood of pathogen transmission. Multiple studies have shown that feral felids are susceptible to contracting diseases from contaminated surfaces or asymptomatic pathogen carriers. However, the etiology of these instances of contagiousness remains obscure [43].

IV. CONCLUSION

It has been noted that felines can display clinical indications suggestive of SARS-CoV-2 infection when their owners, who have been verified to be suffering from COVID-19, undergo laboratory analysis. Unfortunately, the scarcity of examples in this study diminishes its usefulness to some extent. Notwithstanding the elevated incidence of COVID-19 cases in the city of Bandung, Indonesia, the acquisition of samples for diagnostic testing has presented a formidable obstacle. Moreover, it has been observed that certain individuals affected by the COVID-19 disease, whose diagnosis was confirmed by a laboratory, have chosen not to reveal their condition to the relevant authorities owing to the unfavorable stigma linked with the ailment. As a senior professor in environmental health, it is imperative to emphasize the significance of transparency and timely reporting of infectious diseases to prevent further transmission and safeguard public health. It is of utmost importance for individuals who are custodians of *Felis catus* to exercise prudence, given the proclivity of these domesticated quadrupeds to roam freely and cohabit close to *Homo sapiens*. It has been empirically established that the implementation of efficacious hygiene practices, the enforcement of social distancing measures, the maintenance of physical separation, and the adherence to quarantine protocols are all efficacious in mitigating the risk of transmission. The incorporation of whirlpools, footbaths, disinfectants, and hygienic walls necessitates careful deliberation.

Based on the existing corpus of knowledge, it seems highly unlikely that zoonotic transmission of SARS-CoV-2 will transpire. It is important to consider that the possibility of transmission occurring bidirectionally cannot be disregarded. This observation is in line with the findings previously reported by scholars [16]. Through the analysis of serological, molecular, and trial inoculation studies, it has been determined that the SARS-CoV-2 virus possesses zoonotic potential in select animal species. The conclusive demonstration of the transmission of SARS-CoV-2 between animals and humans or vice versa remains uncertain. In light of the present global scenario, it is imperative to adopt a comprehensive strategy to ensure the protection of domesticated animals and captive wildlife from the persistent COVID-19 outbreak. In light of the ongoing COVID-19 pandemic, it is imperative that we extend the same level of care and attention to our animal companions as we do to our human family members in order to mitigate the spread of this highly contagious virus. It is of utmost importance to limit

interpersonal contact with individuals outside the boundaries of one's dwelling and to confine domesticated felines indoors in order to prevent the potential for other companion animals to stray. Adorning our animal companions with masks is ill-advised as it may potentially jeopardize their welfare. It is of utmost importance that individuals who are suffering from a communicable disease abstain from any form of physical interaction with domesticated animals. It is of utmost importance to abstain from participating in behaviors such as osculation, embrace, or sharing a sleeping space with non-human animals, among other similar activities. In the event of an individual contracting SARS-CoV-2, it is recommended that they seek other household members' aid in caring for their pets. If the former alternative is not practicable, it is recommended that individuals wear masks while interacting with their animal companions. It is of utmost importance that the owner takes necessary measures to ensure that their animal companion is provided with nutritionally appropriate sustenance. It is important that companion animals displaying signs of illness or potential COVID-19 infection be expeditiously transported to a veterinary facility for prompt evaluation and management.

ACKNOWLEDGMENT

We thank the Academic Leadership Grant (ALG) for funding this research and the Directorate of Research and Community Service of Universitas Padjadjaran for funding the article processing.

REFERENCES

- [1] Tregoning J S, Brown E S, Cheeseman H M, Flight, K E, Higham S L, Lemm N M, Pierce B F, Stirling D C, Wang Z and Pollock K M 2020 Vaccines for COVID-19 *Clin. Exp. Immunol* 202 162–192 doi:10.1111/cei.13517.
- [2] Kumar S, Nyodu R, Maurya V K and Saxena S K. 2020 Morphology, Genome Organization, Replication, and Pathogenesis of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2). *Coronavirus Disease 2019 (COVID-19)* 23–31 doi:10.1007/978-981-15-4814-7_3.
- [3] Pal M, Berhanu G, Desalegn C and Kandi V 2020 Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-CoV-2): An Update *Cureus* 12 e7423, doi:10.7759/cureus.7423.
- [4] Schoeman D and Fielding B C 2019 Coronavirus Envelope Protein: Current Knowledge *Virology Journal* 16 69 doi:10.1186/s12985-019-1182-0.
- [5] Ren L L, Wang Y M, Wu Z Q, Xiang Z C, Guo L, Xu T, Jiang Y Z, Xiong Y, Li YJ, Li X W, et al 2020 Identification of a Novel Coronavirus Causing Severe Pneumonia in Human: A Descriptive Study *Chinese Medical Journal*. 133 1015–1024 doi:10.1097/CM9.0000000000000722.
- [6] Margolin E, Burgers W A, Sturrock E D, Mendelson M, Chapman R, Douglass N, Williamson A L and Rybicki E P. Prospects for SARS-CoV-2 Diagnostics, Therapeutics and Vaccines in Africa. *Nat Rev Microbiol* 18 690–704 doi:10.1038/s41579-020-00441-3.
- [7] Liu D X, Liang J Q and Fung T S 2021 Human Coronavirus-229E, -OC43, -NL63, and -HKU1 (*Coronaviridae*). *Encyclopedia of Virology* 428–440, doi:10.1016/B978-0-12-809633-8.21501-X.
- [8] Zhu Z, Lian X, Su X, Wu W, Marraro G A and Zeng Y 2020 From SARS and MERS to COVID-19: A Brief Summary and Comparison of Severe Acute Respiratory Infections Caused by Three Highly Pathogenic Human Coronaviruses. *Respiratory Research* 21 224 doi:10.1186/s12931-020-01479-w.
- [9] Baviskar T, Raut D and Bhatt L K 2021 Deciphering Vaccines for COVID-19: Where Do We Stand Today? *Immunopharmacol Immunotoxicol* 43 8–21 doi:10.1080/08923973.2020.1837867.
- [10] BCC Admin Portal Bandung Update statistik covid-19 di Kota Bandung per tanggal 26 April 2023 <https://www.bandung.go.id/information/read/367/update-statistik-covid-19-di-kota-bandung-per-tanggal-26-april-2023> accessed on 15 Mei 2023.
- [11] Dinnata R Y W 2021 Angka Kepemilikan Kucing Naik Signifikan <https://www.ayoyogya.com/explore/pr-39469989/Angka-Kepemilikan-Kucing-di-Indonesia-Naik-Signifikan> Accessed on 15 Mei 2023.
- [12] Salajegheh Tazerji S, Magalhães Duarte P, Rahimi P, Shahabinejad F, Dhakal S, Singh Malik Y, Shehata A A, Lama J, Klein J, Safdar M, et al. 2020 Transmission of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) to Animals: An Updated Review. *Journal of Translational Medicine* 18 358 doi:10.1186/s12967-020-02534-2.
- [13] Giraldo-Ramirez S, Rendon-Marin S, Jaimes J A, Martinez-Gutierrez M, and Ruiz-Saenz J 2021 SARS-CoV-2 Clinical Outcome in Domestic and Wild Cats: A Systematic Review. *Animals* 11 2056, doi:10.3390/ani11072056.
- [14] Zhao J, Cui W, and Tian B. 2020 The Potential Intermediate Hosts for SARS-CoV-2. *Frontiers in Microbiology* 11 2400, doi:10.3389/fmicb.2020.580137.
- [15] Zheng J 2020 SARS-CoV-2: An Emerging Coronavirus That Causes a Global Threat. *Int J Biol Sci* 16 1678–1685 doi:10.7150/ijbs.45053.
- [16] Gryseels S, De Bruyn L, Gyselings R, Calvignac-Spencer S, Leendertz F H and Leirs H 2021 Risk of Human-to-Wildlife Transmission of SARS-CoV-2 *Mammal Review* 51 272–292, doi:10.1111/mam.12225.
- [17] Chiochetti R, Galiazzo G, Fracassi F, Giancola F and Pietra M 2020 ACE2 Expression in the Cat and the Tiger Gastrointestinal Tracts. *Frontiers in Veterinary Science* 7 514 doi:10.3389/fvets.2020.00514.
- [18] Opriessnig T and Huang Y W 2020 Update on Possible Animal Sources for COVID-19 in Humans. *Xenotransplantation* 27 e12621, doi:10.1111/xen.12621.
- [19] Abdurachman F, Paddock R C and Suhartono M 2021 The Pandemic Has a New Epicenter: Indonesia. *The New York Times* Available online <https://www.nytimes.com/2021/07/17/world/asia/indonesia-covid.html>.
- [20] Combs S 2021 Indonesia is a New COVID-19 Epicenter. The Peak Has yet to Come *National Geographic* 2021 Available online: <https://www.nationalgeographic.com/history/article/indonesia-is-new-covid-epicenter-but-the-peak-has-yet-to-come> (accessed on 17 October 2021).
- [21] COVID Protocol v2.0 2021 [covidprotocols.org](https://6169c1c659a7252fcc324939--covid-protocols-web.netlify.app/en/addendum/ward-design-troubleshooting/) Available online: <https://6169c1c659a7252fcc324939--covid-protocols-web.netlify.app/en/addendum/ward-design-troubleshooting/> (accessed on 17 October 2021).
- [22] Ruiz-Arondo I, Portillo A, Palomar A M, Santibáñez S, Santibáñez P, Cervera C and Oteo J A 2021 Detection of SARS-CoV-2 in Pets Living with COVID-19 Owners Diagnosed during the COVID-19 Lockdown in Spain: A Case of an Asymptomatic Cat with SARS-CoV-2 in Europe. *Transbound Emerg Dis* 68 973–976 doi:10.1111/tbed.13803.
- [23] Csiszar A, Jakab F, Valencak T G, Lanszki Z, Tóth G E, Kemenesi G, Tarantini S, Fazekas-Pongor V and Ungvari Z 2020 Companion Animals Likely Do Not Spread COVID-19 but May Get Infected Themselves. *GeroScience* 42 1229–1236 doi:10.1007/s11357-020-00248-3.
- [24] Shiba N, Maeda K, Kato H, Mochizuki M and Iwata H 2007 Differentiation of Feline Coronavirus Type I and II Infections by Virus Neutralization Test. *Vet Microbiol* 2007 124 348–352 doi:10.1016/j.vetmic.2007.04.031.
- [25] Zhang Q, Zhang H, Huang K, Yang Y, Hui X, Gao J, He X, Li C, Gong W, Zhang Y, et al. 2020 SARS-CoV-2 Neutralizing Serum Antibodies in Cats: A Serological Investigation *Microbiology* 1-12
- [26] Zhang Q, Zhang H, Gao J, Huang K, Yang Y, Hui X, He X, Li C, Gong W, Zhang Y, et al. 2020 A Serological Survey of SARS-CoV-2 in Cat in Wuhan *Emerg Microbes Infect* 9 2013–2019 doi:10.1080/22221751.2020.1817796.
- [27] Calvet G A, Pereira S A, Ogrzewalska M, Pauvolid-Corrêa A, Resende P C, Tassinari W, Costa A, Keidel L O, Rocha A S B, da Silva M F B, et al. 2021 Investigation of SARS-CoV-2 Infection in Dogs and Cats of Humans Diagnosed with COVID-19 in Rio de Janeiro, Brazil. *PLOS ONE* 16 e0250853, doi:10.1371/journal.pone.0250853.
- [28] Swelum A A, Shafi M E, Albaqami N M, El-Saadony M T, Elsify A, Abdo M, Taha A E, Abdel-Moneim A M E, Al-Gabri N A, Almaiman A A, et al. 2020 COVID-19 in Human, Animal, and Environment: A Review. *Frontiers in Veterinary Science* 7 578 doi:10.3389/fvets.2020.00578.
- [29] Xu X, Chen P, Wang J, Feng J, Zhou H, Li X, Zhong W and Hao P. 2020 Evolution of the Novel Coronavirus from the Ongoing Wuhan

- Outbreak and Modeling of Its Spike Protein for Risk of Human Transmission. *Sci. China Life Sci.* 63 457–460 doi:10.1007/s11427-020-1637-5.
- [30] Zhou P, Yang X, Wang X G, Hu B, Zhang, L, Zhang W, Si H R, Zhu Y, Li B, Huang C L, et al. 2020 A Pneumonia Outbreak Associated with a New Coronavirus of Probable Bat Origin. *Nature* 579 270–273 doi:10.1038/s41586-020-2012-7.
- [31] Paltrinieri S, Giordano A, Stranieri A and Lauzi S. Feline Infectious Peritonitis (FIP) and Coronavirus Disease 19 (COVID-19): Are They Similar? *Transbound. Emerg. Dis.* 68 1786–1799 doi:10.1111/tbed.13856.
- [32] Delaplace M, Huet H, Gambino A and Le Poder S. 2021 Feline Coronavirus Antivirals: A Review *Pathogens* 10 1150 doi:10.3390/pathogens10091150.
- [33] Rodrigues J P G L M, Barrera-Vilarmau S, Teixeira J M C, Sorokina M, Seckel E, Kastritis P L and Levitt M. 2020 Insights on Cross-Species Transmission of SARS-CoV-2 from Structural Modeling *PLOS Computational Biology* 16, e1008449, doi:10.1371/journal.pcbi.1008449.
- [34] Zhai X, Sun J, Yan Z, Zhang J, Zhao J, Zhao Z, Gao Q, He W T, Veit M and Su S. 2020 Comparison of Severe Acute Respiratory Syndrome Coronavirus 2 Spike Protein Binding to ACE2 Receptors from Human, Pets, Farm Animals, and Putative Intermediate Hosts *J Virol* 94 e00831-20, doi:10.1128/JVI.00831-20.
- [35] Shi J, Wen Z, Zhong G, Yang H, Wang C, Huang B, Liu R, He X, Shuai L, Sun Z, et al. 2020 Susceptibility of Ferrets, Cats, Dogs, and Other Domesticated Animals to SARS-Coronavirus 2 *Science* 368 1016–1020 doi:10.1126/science.abb7015.
- [36] Shi J, Wen Z, Zhong G, Yang H, Wang C, Huang B and Bu Z. 2020 Susceptibility of ferrets, cats, dogs, and other domesticated animals to SARS–coronavirus 2 *Science* 368 (6494) 1016-1020.
- [37] Sun K, Gu L, Ma L and Duan Y. 2020 Atlas of ACE2 gene expression reveals novel insights into transmission of SARS-CoV-2 *Heliyon* 7(1): e05850.
- [38] Sailleau C, Dumarest M, Vanhomwegen J, Delaplace M, Caro V, Kwasiborski A 2020 First detection and genome sequencing of SARS-CoV-2 in an infected cat in France. *Transbound Emerg. Dis.* 67(6) 2324-2328.
- [39] Dhama K, Patel S K, Sharun K, Pathak M, Tiwari R, Yatoo M I, Malik Y S, Sah R, Rabaan A A, Panwar P K, et al. 2020 SARS-CoV-2 Jumping the Species Barrier: Zoonotic Lessons from SARS, MERS and Recent Advances to Combat This Pandemic Virus *Travel Med Infect Dis* 37 101830 doi:10.1016/j.tmaid.2020.101830.
- [40] Villar M, de Mera I G F, Artigas-Jerónimo S, Contreras M, Gortázar C and de la Fuente J. 2020 Coronavirus in Cat Flea: Findings and Questions Regarding COVID-19 *Parasites & Vectors* 13 409 doi:10.1186/s13071-020-04292-y.
- [41] Subbarao K and Mahanty S 2020 Respiratory Virus Infections: Understanding COVID-19 *Immunity* 52 905–909 doi:10.1016/j.immuni.2020.05.004.
- [42] Neira V, Brito B, Agüero B, Berrios F, Valdés V, Gutierrez A, Ariyama N Espinoza P, Retamal P, Holmes E C, et al. 2021 A Household Case Evidences Shorter Shedding of SARS-CoV-2 in Naturally Infected Cats Compared to Their Human Owners *Emerg Microbes Infect* 10 376–383, doi:10.1080/22221751.2020.1863132.
- [43] Musso N, Costantino A, La Spina S, Finocchiaro A, Andronico F and Stracquadanio S 2020 New SARS-CoV-2 Infection Detected in an Italian Pet Cat by RT-qPCR from Deep Pharyngeal Swab *Pathogens* 9(9) 746.