













- [45] Truong, V. D., Mcfeeters, R. F., Thompson, R. T., Dean, L. L., Shofran, B., "Phenolic acid content and composition in leaves and roots of common commercial sweet potato (*Ipomea batatas* L.) cultivars in the United States", *Journal of Food Science*, vol. 72, pp. 343-349, 2007.
- [46] Bellail, A. A., Shaltout, O. E., Youssef, M. M., El Gamal, A. M. A., "Effect of home-cooking methods on phenolic composition and antioxidant activity of sweet potato (*Ipomoea batatas* (L.) Lam.) cultivars grown in Egypt", *Food and Nutrition Science*, vol. 3, pp. 490-499, 2012.
- [47] Xu, B., & Chang, S. K., "Total phenolic, phenolic acid, anthocyanin, Flavan-3-ol, and flavonol profiles and antioxidant properties of pinto and black beans (*Phaseolus vulgaris* L.) as affected by thermal processing", *Journal of Agricultural and Food Chemistry*, vol. 57(11), pp. 4754-4764, 2009.
- [48] Zaro, M. J., Ortiz, L. C., Keunchkarian, S., Chaves, A. R., Vicente, A. R., Concellón, A., "Chlorogenic acid retention in white and purple eggplant after processing and cooking", *LWT - Food Science and Technology*, vol. 64, pp. 802-808, 2015.