























- [39] Ronald A. Holser. 2012. Principal Component Analysis of Phenolic Acid Spectra. *ISRN Spectroscopy*, vol. 2012, Article ID 493203, 5 pages, doi:10.5402/2012/493203.
- [40] Duarte, L.F., Barros, A., Delgado, I., Almeida, C., and Gil, A.M. 2002. Application of FTIR Spectroscopy for the Quantification of Sugars in Mango Juice as a Function of Ripening. *J. Agric. Food Chem.*, 50 (11), pp 3104–3111. DOI: 10.1021/jf011575y.
- [41] F.J. Rambla, S. Garrigues, and M. de la Guardia. 1997. PLS-NIR determination of total sugar, glucose, fructose, and sucrose in aqueous solutions of fruit juices. *Analytica Chimica Acta* 344(1–2):41-53.
- [42] Lijuan Xie, L., Ye, X., and Liu, D. 2009. Quantification of glucose, fructose and sucrose in bayberry juice by NIR and PLS. *Food Chemistry* 114(3):1135-1140.
- [43] Mohsenin, N.N., 1986. *Physical Properties of Plant and Animal Materials*. Gordon and Breach Science Publishers, New York.
- [44] FAO. 1999. CODEX STAN 212-1999 pp.1-5.
- [45] Williamson D. 2004. Study: high-fructose corn sweeteners partly responsible for the obesity epidemic. UNC Gillings School of Global Public Health. USA.
- [46] Hall, K.D. 2008. What is the Required Energy Deficit per unit Weight Loss?. *Int. J. Obes. (Lond)*. 32(3): 573–576. doi: 10.1038/sj.ijo.0803720.
- [47] Makky M., Santosa, Putri R.E., and Nakano K. 2018. Nondestructive evaluation of Bakwan paddy grains moisture content by means of spectrophotometry. *Journal of Physics: Conference Series*. 985:012012. doi :10.1088/1742-6596/985/1/012012