

- farmland in China using time series MODIS NDVI,” *Sci. Total Environ.*, vol. 755, 2021, doi: 10.1016/j.scitotenv.2020.142651.
- [13] R. Moreno, N. Ojeda, J. Azócar, C. Venegas, and L. Inostroza, “Application of NDVI for identify potentiality of the urban forest for the design of a green corridors system in intermediary cities of Latin America: Case study, Temuco, Chile,” *Urban For. Urban Green.*, vol. 55, no. 126821, 2020, doi: 10.1016/j.ufug.2020.126821.
- [14] S. Testa, K. Soudani, L. Boschetti, and E. B. Mondino, “MODIS-derived EVI, NDVI and WDRVI time series to estimate phenological metrics in French deciduous forests,” *Int J Appl Earth Obs Geoinf.*, vol. 64, no. July 2017, pp. 132–144, 2018, doi: 10.1016/j.jag.2017.08.006.
- [15] F. F. Gerard, C. T. George, G. Hayman, C. Chavana-Bryant, and G. P. Weedon, “Leaf phenology amplitude derived from MODIS NDVI and EVI: Maps of leaf phenology synchrony for Meso- and South America,” *Geosci. Data J.*, vol. 00, pp. 1–14, 2020, doi: 10.1002/gdj3.87.
- [16] P. Karkauskaite, T. Tagesson, and R. Fensholt, “Evaluation of the Plant Phenology Index (PPI), NDVI and EVI for Start-of-Season Trend Analysis of the Northern Hemisphere Boreal Zone,” *Remote Sens.*, vol. 9, no. 485, pp. 21–21, 2017, doi: 10.3390/rs9050485.
- [17] M. Gandhi G, S. Parthiban, N. Thummalu, and C. A, “NdvI : Vegetation change detection using remote sensing and gis – A case study of Vellore District,” *Procedia - Procedia Comput. Sci.*, vol. 57, pp. 1199–1210, 2015, doi: 10.1016/j.procs.2015.07.415.
- [18] R. Colombo, D. Bellingeri, D. Fasolini, and C. M. Marino, “Retrieval of leaf area index in different vegetation types using high resolution satellite data,” *Remote Sens. Environ.*, vol. 86, pp. 120–131, 2003, doi: 10.1016/S0034-4257(03)00094-4.
- [19] A. Kross, H. McNairn, D. Lapen, M. Sunohara, and C. Champagne, “Assessment of RapidEye vegetation indices for estimation of leaf area index and biomass in corn and soybean crops,” *Int. J. Appl. Earth Obs. Geoinf.*, vol. 34, pp. 235–248, 2015, doi: 10.1016/j.jag.2014.08.002.
- [20] L. Wang *et al.*, “Effects of growth stage development on paddy rice leaf area index prediction models,” *Remote Sens.*, vol. 11, no. 361, pp. 1–18, 2019, doi: 10.3390/rs11030361.
- [21] J. Zhao *et al.*, “Estimating fractional vegetation cover from leaf area index and clumping index based on the gap probability theory,” *Int. J. Appl. Earth Obs. Geoinf.*, vol. 90, no. 102112, 2020, doi: 10.1016/j.jag.2020.102112.
- [22] Q. Wang, S. Adiku, J. Tenhunen, and A. Granier, “On the relationship of NDVI with leaf area index in a deciduous forest site,” *Remote Sens. Environ.*, vol. 94, pp. 244–255, 2005, doi: 10.1016/j.rse.2004.10.006.
- [23] Suwarsono *et al.*, “Pengembangan metode penentuan Indeks Luas Daun pada penutup lahan hutan dari data satelit penginderaan jauh SPOT-2,” *J. Penginderaan Jauh*, vol. 8, pp. 50–59, 2011.
- [24] R. Darvishzadeh *et al.*, “Analysis of Sentinel-2 and rapidEye for retrieval of leaf area index in a saltmarsh using a radiative transfer model,” *Remote Sens.*, vol. 11, no. 671, 2019, doi: 10.3390/rs11060671.
- [25] T. Mannschatz, B. P. E. Borg, K. Feger, and P. Dietrich, “Uncertainties of LAI estimation from satellite imaging due to atmospheric correction,” *Remote Sens. Environ.*, vol. 153, pp. 24–39, 2014, doi: 10.1016/j.rse.2014.07.020.
- [26] L. A. Zadeh, “Fuzzy Sets,” *Inf. Control*, vol. 353, pp. 338–353, 1965.
- [27] D. Saadoud, M. Hassani, F. José, and M. Peinado, “Application of fuzzy logic approach for wind erosion hazard mapping in Laghouat region (Algeria) using remote sensing and GIS,” *Aeolian Res.*, vol. 32, no. February, pp. 24–34, 2018, doi: 10.1016/j.aeolia.2018.01.002.
- [28] S. Sarkar, S. M. Parihar, and A. Dutta, “Environmental Modelling & Software Fuzzy risk assessment modelling of East Kolkata Wetland Area : A remote sensing and GIS based approach,” *Environ. Model. Softw.*, vol. 75, pp. 105–118, 2016, doi: 10.1016/j.envsoft.2015.10.003.
- [29] G. M. Foody, “Fuzzy modelling of vegetation from remotely sensed imagery,” *Ecol. Modell.*, vol. 85, pp. 3–12, 1996.
- [30] T. Semeraro, G. Mastroleo, A. Pomes, A. Luvisi, E. Gissi, and R. Aretano, “Modelling fuzzy combination of remote sensing vegetation index for durum wheat crop analysis,” *Comput. Electron. Agric.*, vol. 156, no. December 2018, pp. 684–692, 2019, doi: 10.1016/j.compag.2018.12.027.
- [31] Y. Zhang, Q. Li, X. Du, and H. Wang, “Spatially explicit mapping of phenological transition zones: A fuzzy-logic approach,” *Agric. For. Meteorol.*, vol. 295, no. 108201, 2020, doi: 10.1016/j.agrformet.2020.108201.
- [32] R. M. Gonçalves, A. Saleem, H. A. A. Queiroz, and J. L. Awange, “A fuzzy model integrating shoreline changes, NDVI and settlement influences for coastal zone human impact classification,” *Appl. Geogr.*, vol. 113, no. 102093, 2019, doi: 10.1016/j.apgeog.2019.102093.
- [33] C. M. Rulinda, A. Dilo, W. Bijker, and A. Stein, “Characterising and quantifying vegetative drought in East Africa using fuzzy modelling and NDVI data,” *J. Arid Environ.*, vol. 78, pp. 169–178, 2012, doi: 10.1016/j.jaridenv.2011.11.016.
- [34] S. Ghosh, A. Das, T. K. Hembram, S. Saha, B. Pradhan, and A. M. Alamri, “Impact of COVID-19 induced lockdown on environmental quality in four Indian megacities Using Landsat 8 OLI and TIRS-derived data and Mamdani fuzzy logic modelling approach,” *Sustain.*, vol. 12, no. 5464, pp. 1–24, 2020, doi: 10.3390/su12135464.
- [35] N. M. Sari and D. Kushardono, “Analisis dampak pembangunan infrastruktur Bandara Internasional Jawa Barat terhadap alih fungsi lahan pertanian melalui citra satelit resolusi tinggi,” *J. Geogr.*, vol. 11, no. 2, pp. 146–162, 2019, doi: 10.24114/jg.v11i2.13470.
- [36] J. L. Roujean and F. M. Breon, “Estimating PAR absorbed by vegetation from bidirectional reflectance measurements,” *Remote Sens. Environ.*, vol. 51, no. 3, pp. 375–384, 1995, doi: 10.1016/0034-4257(94)00114-3.