













- agricultural waste," *Int J Appl Sci Biotechnol*, vol. 8, no. 3, pp. 336–342, Sep. 2020. doi: 10.3126/ijasbt.v8i3.31566.
- [14] R. Mitra, T. Xu, G. Q. Chen, H. Xiang, and J. Han, "An updated overview on the regulatory circuits of polyhydroxyalkanoates synthesis," *Microbial Biotechnology*, vol. 15, no. 5, pp. 1446–1470, Sep. 2022. doi: 10.1111/1751-7915.13915.
- [15] B. Gutschmann, M. M. Simoes, T. Schiewe, E. S. Schroter, M. Munzberg, P. Neubauer, A. Bockisch, and S. L. Riedel, "Continuous feeding strategy for polyhydroxyalkanoate production from solid waste animal fat at laboratory- and pilot-scale," *Microb Biotechnol*, vol. 16, no. 2, pp. 295–306, Aug. 2023, doi: 10.1111/1751-7915.14104.
- [16] D. Vicente, D. N. Proença, and P. V. Morais, "The role of bacterial polyhydroxyalkanoate (PHA) in a sustainable future: A review on the biological diversity," *International Journal of Environmental Research and Public Health*, vol. 20, no. 4, p. 2959, Feb. 2023. doi: 10.3390/ijerph20042959.
- [17] W. Zhou, S. Bergsma, D. I. Colpa, G. J. W. Euverink, and J. Krooneman, "Polyhydroxyalkanoates (PHAs) synthesis and degradation by microbes and applications towards a circular economy," *Journal of Environmental Management*, vol. 341, pp. 1–13, Sep. 2023. doi: 10.1016/j.jenvman.2023.118033.
- [18] J. R. Gómez-Cardozo, R. Velasco-Bucheli, N. Marín-Pareja, O. S. Ruiz-Villadiego, G. A. Correa-Londoño, and A. L. Mora-Martínez, "Fed-batch production and characterization of polyhydroxybutyrate by *Bacillus megaterium* LVN01 from residual glycerol," *DYNA (Colombia)*, vol. 87, no. 214, pp. 111–120, May. 2020, doi: 10.15446/dyna.v87n214.83523
- [19] J. Mascarenhas and K. Aruna, "Production and optimization of polyhydroxyalkanoate obtained from *Bacillus megaterium* JHA," *Journal of Applied Biotechnology Reports*, vol. 8, no. 4, pp. 346–360, Dec. 2021. doi: 10.30491/jabr.2020.242493.1263.
- [20] S. Mohammed and L. Ray, "Polyhydroxyalkanoate recovery from newly screened *Bacillus* sp. LPPI-18 using various methods of extraction from Loktak Lake sediment sample," *Journal of Genetic Engineering and Biotechnology*, vol. 20, p. 115, Aug. 2022. doi: 10.1186/s43141-022-00392-7.
- [21] D. H. Vu, S. Wainaina, M. J. Taherzadeh, D. Åkesson, and J. A. Ferreira, "Production of polyhydroxyalkanoates (PHAs) by *Bacillus megaterium* using food waste acidogenic fermentation-derived volatile fatty acids," *Bioengineered*, vol. 12, no. 1, pp. 2480–2498, Jun. 2021. doi: 10.1080/21655979.2021.1935524.
- [22] M. Geethu, H. R. Chandrashekar, and M. S. Divyashree, "Statistical optimisation of polyhydroxyalkanoate production in *Bacillus endophyticus* using sucrose as sole source of carbon," *Arch Microbiol*, vol. 203, no. 10, pp. 5993–6005, Sep. 2021. doi: 10.1007/s00203-021-02554-6.
- [23] N. Haedar, D. Suherman, Z. Dwyana, Heriadi, and M. Masri, "The potency of bioplastic polyhydroxyalkanoate (PHA) producing bacteria isolated from palm oil mill waste," *Elkawnie: Journal of Islamic Science and Technology*, vol. 8, no. 1, pp. 93–107, Jun. 2022. doi: 10.22373/ekw.v8i1.11145.
- [24] H. Javaid, A. Nawaz, N. Riaz, H. Mukhtar, I. Ul-Haq, K. A. Shah, H. Khan, S. M. Nawvi, S. Shakoor, A. Rasool, K. Ullah, R. Manzoor, I. Kaleem, and G. Murtaza, "Valorization of biomass and synthetic waste," *Molecules*, vol. 25, no. 23, p. 5539, Nov. 2020. doi: 10.3390/molecules25235539
- [25] T. Khamkong, W. Penkhruue, and S. Lumyong, "Optimization of production of polyhydroxyalkanoates (PHAs) from newly isolated *Ensifer* sp. Strain HD34 by Response Surface Methodology," *Processes*, vol. 10, no. 8, Aug. 2022. doi: 10.3390/pr10081632.
- [26] D. Li, M. Gao, Y. Qiu, X. Ma, F. Wang, J. Li, and L. Yu, "Strategy for economical and enhanced polyhydroxyalkanoate production from synergistic utilization of palm oil and derived wastewater by activated sludge," *Bioresour Technol*, vol. 370, p. 128581, Feb. 2023, doi: 10.1016/j.biortech.2023.128581
- [27] B. A. Ramsay, K. Lomaliza, C. Chavarie, B. Dube, P. Bataille, and J. A. Ramsay, "Production of poly-( $\beta$ -hydroxybutyric-co- $\beta$ -hydroxyvaleric) acids," vol. 56, no. 7, pp. 2093–2098, Jul. 1990. doi: 10.1128/aem.56.7.2093-2098.1990
- [28] P. J. Senior, G. A. Beech, G. A. F. Ritchie, and E. A. Dawes, "The role of oxygen limitation in the formation of poly-p-hydroxybutyrate during batch and continuous culture of *Azotobacter beijerinckii*," *Biochem J*, vol. 128, no. 5, pp. 1193–1201, Aug. 1972. doi: 10.1042/bj1281193.
- [29] Y. Zhang, W. Sun, H. Wang, and A. Geng, "Polyhydroxybutyrate production from oil palm empty fruit bunch using *Bacillus megaterium* R11," *Bioresour Technol*, vol. 147, pp. 307–314, Nov. 2013, doi: 10.1016/j.biortech.2013.08.029.
- [30] F. Verdini, S. Tabasso, F. Mariatti, F. Bosco, C. Mollea, E. C. Gaudino, A. Cirio, and G. Cravotto, "From agri-food wastes to polyhydroxyalkanoates through a sustainable process," *fermentation*, vol. 8, no. 10, pp. 1–18, Oct. 2022. doi: 10.3390/fermentation8100556.
- [31] A. B. Devi, C. V. Nachiyar, T. Kaviyarasi, and A. V. Samrot, "Characterization of polyhydroxybutyrate synthesized by *Bacillus cereus*," *Int J Pharm Pharm Sci*, vol. 7, no. 3, pp. 140–144, Jan. 2015. ISSN-0975-1491
- [32] F. Crisafi, F. Valentino, F. Micolucci, and R. Denaro, "From organic wastes and hydrocarbons pollutants to polyhydroxyalkanoates: bioconversion by terrestrial and marine bacteria," *Sustainability*, vol. 14, no. 14, pp. 1–29, Jul. 2022. doi: 10.3390/su14148241
- [33] A. T. Ubando, C. B. Felix, and W. H. Chen, "Biorefineries in circular bioeconomy: A comprehensive review," *Bioresour Technol*, vol. 299, p. 122585, Mar. 2020. doi: 10.1016/j.biortech.2019.122585.
- [34] E. Gesteiro, L. Guijarro, F. J. Sanchez-Muniz, M. C. Vidal-Carou, A. Troncoso, L. Venanci, Vicente Jimeno, J. Quilez, A. Anadon, and M. Gonzalez-Gross, "Palm oil on the edge," *Nutrients*, vol. 11, no. 9, pp. 1–36, Aug. 2019. doi: 10.3390/nu11092008
- [35] S. Shah and A. Kumar, "Production and characterization of polyhydroxyalkanoates from industrial waste using soil bacterial isolates," *Braz J Microbiol*, vol. 52, no. 2, pp. 715–726, Jun. 2021. doi: 10.1007/s42770-021-00452-z/Published.
- [36] J. H. C. Chin, M. R. Samian, and Y. M. Normi, "Characterization of polyhydroxyalkanoate production capacity, composition and weight synthesized by *Burkholderia cepacia* JC-1 from various carbon sources," *Heliyon*, vol. 8, no. 3, pp. 1–11, Mar. 2022. doi: 10.1016/j.heliyon.2022.e09174.
- [37] T. Sriyapai, T. Chuarung, K. Kimbara, S. Samosorn, and P. Sriyapai, "Production and Optimization of polyhydroxyalkanoates (PHAs) from *Paraburkholderia* sp. PFN 29 under submerged fermentation," *Electronic Journal of Biotechnology*, vol. 56, pp. 1–11, Mar. 2022. doi: 10.1016/j.ejbt.2021.12.003.
- [38] N. O. Zhila, K. Y. Sapozhnikova, E. G. Kiselev, A. D. Vasiliev, I. V. Nemtsev, E. I. Shishatskaya, and T. G. Volova, "Properties of degradable polyhydroxyalkanoates (PHAs) synthesized by a new strain, *Cupriavidus necator* IBP/SFU-1, from various carbon sources," *Polymers (Basel)*, vol. 13, no. 18, pp. 1–19, Sep. 2021. doi: 10.3390/polym13183142.
- [39] D. Correa-Galeote, L. Argiz, A. Val del Rio, A. Mosquera-Corral, B. Juarez-Jimenez, J. Gonzales-Lopez, and B. Rodelas, "Dynamics of PHA-accumulating bacterial communities fed with lipid-rich liquid effluents from fish-canning industries," *Polymers (Basel)*, vol. 14, no. 7, pp. 1–22, Mar. 2022. doi: 10.3390/polym14071396.
- [40] R. Sohail, N. Jamil, I. Ali, and S. Munir, "Animal fat and glycerol bioconversion to polyhydroxyalkanoate by produced water bacteria," *e-Polymers*, vol. 20, pp. 92–102, Mar. 2020. doi: 10.1515/epoly-2020-0011
- [41] D. M. Miu, M. C. Eremia, and M. Moscovici, "Polyhydroxyalkanoates (PHAs) as biomaterials in tissue engineering: production, isolation, characterization," *Materials*, vol. 15, no. 4, Feb. 2022. doi: 10.3390/ma15041410.
- [42] N. O. Zhila, K. Y. Sapozhnikova, E. G. Kiselev, E. I. Shishatskaya, and T. G. Volova, "Synthesis and properties of polyhydroxyalkanoates on waste fish oil from the production of canned sprats," *Processes*, vol. 11, no. 7, Jul. 2023. doi: 10.3390/pr11072113.
- [43] M. Ishii-Hyakutake, S. Mizuno, and T. Tsuge, "Biosynthesis and characteristics of aromatic polyhydroxyalkanoates," *Polymers (Basel)*, vol. 10, no. 11, pp. 1–24, Nov. 2018. doi: 10.3390/polym10111267.
- [44] P. Tanikkul, G. L. Sullivan, S. Sarp, and N. Pisutpaisal, "Biosynthesis of medium chain length polyhydroxyalkanoates (mcl-PHAs) from palm oil," *Case Studies in Chemical and Environmental Engineering*, vol. 2, no. 100045, p. 1–5, Sep. 2020. doi: 10.1016/j.cscee.2020.100045.